



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

July 1, 2015

Debbie Ziehr
Regulatory Administrator
The Andersons Lawn and Fertilizer Division, Inc.
The Andersons Inc.
P. O. Box 119
Maumee, OH 43537

Subject: Notification per PRN 98-10 – Addition of nitrogen and phosphorus restrictions
Product Name: The Andersons Dimension Herbicide III
EPA Registration Number: 9198-121
Application Date: June 3, 2015
Decision Number: 503596

Dear Ms. Ziehr:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

If you have any questions, you may contact Deirdre Sunderland at (703) 603-0851 or via email at sunderland.deirdre@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague". The signature is written in a cursive style with a long horizontal flourish extending to the right.

Kathryn V. Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

THE ANDERSONS DIMENSION HERBICIDE III

This product is a herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs.

Optional marketing claim:



Dispersing Granule Technology

www.andersonsgolfproducts.com

Andersons Golf Products and Contec DG are registered trademarks of The Andersons Inc.

- Utilizes DG^{Pro}® dispersible carrier.
- Contains Nutrisphere-N[®]

NOTIFICATION

9198-121

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

07/01/2015

ACTIVE INGREDIENTS:

Dithiopyr, 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S- dimethyl ester.....	0.164%
---	--------

INERT INGREDIENTS	<u>99.836%</u>
--------------------------------	----------------

TOTAL.....	100.00%
------------	---------

**KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS**

EPA Reg. No. 9198-121

EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
(Underlined letter is first letter used in run code on end of bag)

NET WEIGHT _____ LBS.

The Andersons Dimension Herbicide III

GUARANTEED ANALYSIS

Total Nitrogen (N).....	16%
Available Phosphate (P ₂ O ₅).....	4%
Soluble Potash (K ₂ O).....	4%
Primary Plant Food derived from: Urea, Muriate of Potash, Ammoniated Phosphates.	

For products being sold into Florida:

Florida Applications:

We recommend that you follow the Florida Green Industries BMP's at:

http://www.flaes.org/pdf/BMP_Book_final.pdf

We recommend that you follow the Florida Golf Course BMP's at: <http://www.flaes.org/pdf/glfbmp07.pdf>

Nitrogen Notice: For those states with nitrogen application restrictions, including but not limited to MD, NH, NJ, and VA, do not apply more than x.xx lbs. of this product per 1,000 sq. ft.

Phosphate Notice: This lawn/turf fertilizer contains phosphorous and is only for nonagricultural use on (i) turf during its first growing season, (ii) on turf areas being repaired or renovated, and (iii) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not for routine maintenance of turf.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN

CAUTION

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Read the entire label before using this product.

Use only according to label instructions.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 7.0 pounds per 1000 square feet (equal to 0.5 pound active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes and greens. This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

TURFGRASS SAFETY

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

Cool-Season Grasses

Bentgrass, creeping*
 Agrostis palustris
Bluegrass, Kentucky
 Poa pratensis
Fescue, fine**
 Festuca rubra
Fescue, tall
 Festuca arundinacea
Ryegrass, perennial
 Lolium perenne

Warm-Season Grasses

Bahiagrass
 Paspalum notatum
Bermudagrass
 Cynodon dactylon
Buffalograss***
 Buchloe dactyloides
Carpetgrass
 Axonopus affinis
Centipedegrass
 Eremochloa ophiuroides

Kikuyugrass
 Pennisetum clandestinum
St. Augustinegrass
 Stenotaphrum secundatum
Zoysiagrass
 Zoysia japonica

* Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested.

DO NOT apply this product to Colonial bentgrass (*Agrostis tenuis*) varieties.

**Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.

***DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

NOTE: It is recommended that use of this product be made on a trial basis initially to determine its safety under local use conditions and turf management practices before making widespread use.

RESEEDING, OVERSEEDING, OR SPRIGGING

Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

GOLF COURSE PUTTING GREENS

Putting greens should be well-drained and managed to promote the development of deep rooting systems. The use of this product on greens without good root development may result in undesirable turfgrass injury.

For golf courses which practice overseeding putting greens consisting of seasonal turfgrasses, application of this product to overseeded greens should be made within four weeks or less before the desired transition period from the overseeded turfgrasses back to the established "base" turfgrass. Avoid application of this product to golf course putting greens that are under stress.

Undesirable thinning, yellow and rapid decline of established *Poa annua* may occur under stress conditions. DO NOT use this product on greens that have desirable *Poa annua* cover where summer heat and/or other stresses to *Poa annua* can occur and loss of some or all of that *Poa annua* cover is unacceptable. Use this product where the goal is to control and/or eliminate some or all of the *Poa annua* and where this loss can be tolerated.

Use of this product on 'Tifgreen' (328) hybrid bermudagrass greens may result in injury. Prior to use, test on a small section of turf.

DO NOT apply this product less than 60 days before covering greens for winter protection.

OTHER USE PRECAUTIONS

The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turfs only. DO NOT apply to flowers, vegetables, shrubs, or trees.

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.

Keep people and pets off treated areas until dust has settled.

APPLICATION DIRECTIONS

**Table 1: Recommended Application Use Rates*
(lbs of this product/1000 sq. ft.)**

PROGRAM	1	2	3	4
USE (turfgrass cut)	Preemergence (high)	Preemergence (low) Postemergence (high)	Preemergence (low) Postemergence (low)	Golf Course Putting Green Mowing Height less than 1/4 inch**
REGION				
North	2.3	2.6	3.5	3.5
Transition	2.6	3.5	5.2	3.5
South	3.5	5.2	7.0**	3.5+3.5
Coastal South	5.2	7.0**	7.0**	5.2+3.5+3.5

*DO NOT apply more than 7 lbs. per application, and no more than 21 lbs. per year. Postemergence control is limited; see "Crabgrass Control" section below and "Precautions" section above. Regions include areas listed below.

North: All other areas.

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI.

South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX.

Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

**May require split or sequential applications for full control.

CONTROL OF CRABGRASS

Preemergence and Early Postemergence Control

This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.

It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

Application Frequency and Timing

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall.

DO NOT apply more than 7 lbs. per 1000 sq. ft. per application, and no more than 21 lbs. per 1000 sq. ft. per year.

Spring Applications

For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 above, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use Program 1 for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.

Use Program 2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut high (e.g., homeowner lawns).

Use Program 3 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways) and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut low (e.g., golf fairways).

Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

Fall Applications

This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring.

The fall application should be followed by an appropriate spring application to provide season-long control.

Tips for Improved Control

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.

Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications. Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence:

barnyardgrass
bluegrass (annual, roughstalk)
crowfootgrass
foxtail (yellow & green)*
kikuyugrass
ryegrass (annual & perennial)
smutgrass

lespedeza (common)
oxalis (buttercup, creeping & yellow woodsorrel)
purslane (common)
speedwell (corn)
spurge (prostrate & spotted)
goosegrass

*Also controlled at the Program 2 rates.

Fall Applications. Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence:

bluegrass (annual)-Poa annua
bittercress
chickweed
henbit

geranium (Carolina)
parsley-piert
pineappleweed
shepherdspurge

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

Suggested Spreader Settings
(Available at final printing)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. **STORAGE:** Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If not available, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION - LANDSCAPE ORNAMENTALS

This product provides preemergence control of listed annual grass and broadleaf weeds in landscape ornamentals. It may be applied to soil surfaces where ornamental plants are grown for aesthetic purposes in ornamental gardens, and residential properties.

Ornamental plants on this label have shown tolerance to this product when applied as directed. It is impossible, however, to evaluate tolerance for all ornamental plant species or varieties or under all possible growing conditions. Under your growing conditions, a limited area involving only a few plants should be treated for tolerance evaluation prior to large-scale applications.

USE RATES

Apply this product with a properly calibrated spreader which will assure accurate, even particle distribution. This product may be applied as a single application or as a split application. Refer to the following table for product recommendations when treating a small area.

DO NOT apply more than 7 pounds per 1,000 square feet per application and no more than 21 pounds (11 pounds in New York) per 1,000 square feet per year. Do not apply in Nassau and Suffolk Counties of New York.

Pounds of Product
per 1,000 square feet
7

Ounces of Product
per 100 Square Feet
11.2

APPLICATION TIMING AND RECOMMENDATIONS

This product is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination or to bare ground. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. Once an application of this product has been made, do not disturb the soil surface as the herbicide barrier will be broken.

PRECAUTIONS

- Only apply this product to established ornamentals.
- Do not apply this product directly to bare root of ornamental plants as injury may result.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply to soil around ornamental plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes or disease.
- To avoid foliar damage to ornamentals, irrigate immediately after application to remove herbicide particles from the foliage.
- Do not apply under conditions that would affect product distribution. Wind could affect distribution and move product onto sensitive species. Uneven product distribution will result in uneven weed control.
- Do not apply this product on grasses grown for seed.
- Do not graze livestock or feed forage cut from areas treated with this product.

TOLERANT ORNAMENTALS

When applied as directed under the conditions described on this label, ornamentals listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing condition. Follow directions given above to determine plant tolerance under your growing conditions prior to large scale use.

COMMON NAME	BOTANICAL NAME	TOLERANT CULTIVARS
Abelia, Dwarf	<i>Abelia X Grandiflora</i>	Nana
Ajuga	<i>Ajuga reptans</i> <i>Ajunga genevensis</i>	Bronze Bronze Beauty
Almond, Flowering	<i>Prunus gladiosa</i>	
Apple*	<i>Malus pumila</i>	
Arbovitae	<i>Thuja occidentalis</i>	Nigra Pyramidalis Smaragh Techny Woodwardii
Arborvitae, Golden	<i>Thuja orientalis</i>	
Aster, Chinese	<i>Callistephus chinensis</i>	Dwarf Queen
Ash, Green	<i>Fraxinus pennsylvanica</i>	
Ash, Mountain	<i>Sorbus acucparia</i>	
Ash, Purple	<i>Fraxinus americana</i>	
Azalea	<i>Rhododendron spp.</i>	Brillant Buccaneer Carror Chimes (Belgian) Exbury Fashion Hardijzer Beauty Hershey Red Higasa Hinocrimson Holland (Hybrid) Marion Lee Northern Lights Orange Cup Orchid Lights Snow

		Southern Charm
Azalea, Flame	<i>Rhododendron calendulaceum</i>	
Azalea, Kirishima		
Bamboo, Heavenly		
Barberry	<i>Berberis thunbergii</i>	Aurea Dwarf Pigmy Green Kobold Pygmy Red Rose Glow
Barberry, Purple		Atropurpurea
Bearberry (common)	<i>Arctostaphylos uva-ursti</i>	Massachusetts
Birch, River	<i>Betula nigra</i>	
Blueberry*	<i>Vaccinium spp.</i>	Bluecorp Blue Jay Jersey North Blue Northland
Bottlebrush	<i>Callistemon citrinum</i>	
Boxwood, Japanese		Japonica
Boxwood, Weller	<i>Buxus sempervirens</i>	
Broom	<i>Cytisus spp.</i>	Moonlight
Broom	<i>Genista pilosa</i>	Vancouver Gold
Bugle Carpet		
Carex, Variegated	<i>Carex</i>	
Camellia	<i>Camellia japonica</i> <i>Camellia sasangua</i>	Debutante Mathotiana Supreme Chansonette
Cedar, Red	<i>Juniperus virginiana</i>	
Cockscomb, Plumosa	<i>Celosia cristata</i>	Red Kewpie
Coleus	<i>Coleus blumei</i>	
Copper leaf	<i>Acalypha wilkesiana</i>	
Cotoneaster	<i>Cotonester apiculatus</i>	
Cycads	<i>Cycads revoluta</i>	
Coyotebrush	<i>Baccharis pelularis</i>	
Cyrpress, Bald	<i>Taxodium distichum</i>	
Cypress, Italian	<i>Cupressus sempervirens</i>	Glauca
Cypress, Leyland	<i>Cupressocyparis leylandii</i>	
Cypress, Japanese False	<i>Chamaecyparis obtusa</i>	Gracilis
Daffodil	<i>Narcissus spp.</i>	King Alfred
Daylilly	<i>Heemerocallis spp.</i>	Aztec Gold Bright Yellow (Hybrid) Single Gold (Evergreen) Wilson's Yellow
Dogwood	<i>Cornus florida</i>	
Dogwood, American	<i>Cornus sericea</i>	
Douglas Fir	<i>Pseudotsuga menziesii</i>	
Dusty Miller	<i>Senecio cineraria</i>	
Elm, Drake	<i>Ulmus parvifolia</i>	
Euonymus	<i>Euonymus fortunei</i>	Argenteo-variegata Auereo-marginata Colorata

Impatiens	<i>Impatiens wallerana</i>	
Iris	<i>Iris spp.</i>	Wedgewood Dwarf blue
Ivy, English	<i>Hedera helix</i>	Bulgaria Thorndale
Jasmine, Asian	<i>Trachelopermum asiaticum</i>	
Juniper	<i>Juniperus spp.</i>	Arcadia Armstrong Bar Harbor
	<i>Juniperus horizontalis</i>	Blue Chip Blue Pacific Blue Rug
	<i>Juniperus horizontalis</i>	Blue Star Broadmoor Buffalo Calgary Carpet Emerald Sea Emerald Spreader Endora Compacta Fuitlandi Green Gold Tip Hetzi Hughes Manhattan Blue Parsoni Pfitzeriana Plumosa Prince of Wales Procumbens Dwarf San Jose Sargent Blude Sargent Green Scandia Scopulorum Moonglow Scopulorum Skyrocket Spartan
	<i>Juniperus chinensis</i>	Tamariscifolia
	<i>Juniperus sabina</i>	Weberi Youngstown Yukon Belle
King Palm	<i>Archontophoenix cunninghamiana</i>	
Laurel, Australian	<i>Pittosporum tobira</i>	
Laurel, Mountain	<i>Kalmia latifolia</i>	
Leucothoe	<i>Leucothoe fontanesiana</i>	
Lily, Africian	<i>Agapanthus africanus</i>	Albus Peter Pan
Lily, African Blue		
Lily of the Valley	<i>Pieris japonica</i>	Mt. Fire
Lilyturf	<i>Liriope muscari</i>	Evergreen Giant Lilac Beauty Majestic Monroe White

		Variegata
Ligustrum, Japanese	<i>Ligustrum japonicum</i>	
Liriope, Green	<i>Liriope spicata</i>	
Maple, Japanese	<i>Acer japonicum</i>	
Maple, Norway	<i>Acer platanoides</i>	
Maple, Red	<i>Acer rubrum</i>	
Maple, Sivler	<i>Acer saccharinum</i>	
Maple, Sugar	<i>Acer saccharum</i>	
Marigold	<i>Tagetes patula</i>	Honeycomb Wheeler's Dwarf Variegata
Mock Orange*	<i>Philadelphus spp</i>	Golden Snowflake Double White
Mopondo Grass	<i>Ophiopogon japonicus</i>	
Moss Rose	<i>Portulaca grandiflora</i>	Sunnyside
Myrtle, Crape	<i>Lagerstoemia indica</i>	Faurei Langer Muskogee Standard Pink
Myrtle, Wax	<i>Myrica cerifer</i>	
Nadina	<i>Nadina domestica</i>	Compacta Nana
Oleander	<i>Nerium oleander</i>	Hardy Red Petite Pink Sister Agnes
Oak, Laurel	<i>Quercus laurifolia</i>	
Oak, Pin	<i>Quercus palustris</i>	
Oak, Red	<i>Quercus rubra</i>	
Oak, Southern L	<i>Quercus virginiana</i>	
Oak, Willow	<i>Quercus phellos</i>	
Oteospermum	<i>Osteospermum fruitcosum</i>	Wirligig
Palm, Bangalow		
Pampas Grass	<i>Cortadaria selloana</i>	
Paper Flower	<i>Bougainvillea glabra</i>	Barbara Karst
Pachysandra	<i>Pachysandra terminalis</i>	
Peach*	<i>Prunus persica</i>	
Periwinkle, Dwarf	<i>Vinca minor</i>	
Petunia	<i>Petunia X hybrida</i>	Picoti
Photinia, Red Tip	<i>Photinia X fraseri</i>	
Pieris	<i>Pieris japonica</i>	
Pine, Australian	<i>Pinus nigra</i>	
Pine, Japanese Black	<i>Pinus thumbergiana</i>	
Pine, Loblolly	<i>Pinus teada</i>	
Pine, Longleaf	<i>Pinus palustris</i>	
Pine, Mugho	<i>Pinus mugho</i>	
Pine, Scotch	<i>Pinus sylvestris</i>	
Pine, Slash	<i>Pinus eliottii</i>	
Pine, Swiss Mt.	<i>Pinus mugo</i>	
Pine, Virginia	<i>Pinus virginiana</i>	
Pine, White	<i>Pinus strobus</i>	
Pineapple, Guava	<i>Feijoa fellowaiana</i>	
Pittosporum, Japan		

Potentilla	<i>Potentilla fruticosa</i>	Abbotswood
Privet	<i>Ligustrum japonicum</i>	Golden Vicary Regal Texanum Yellow Tipped Wax
Privet, Glossy	<i>Ligustrum lucidum</i>	
Pyracantha	<i>Pyracantha koidzumii</i>	
Queen Pam	<i>Arecastrum rammanzoffianum</i>	
Quince, Japanese*		
Rhododendron	<i>Rhododendron spp</i>	Album Cunningham White Fashion Hardy PJM Purple Gem Silvery Pink
Rhododendron, Carolina	<i>Rhododendron carolinianum</i>	
Rhododendron, Catawba	<i>Rhododendron catawbiense</i>	
Rodie Max (Rosebay)	<i>Rhododendron maximum</i>	
Rose	<i>Rosa banksiaae</i>	Luta
Rosemary*	<i>Rosmarinum officinalis</i>	
Rosemary, Bog	<i>Andromeda polifolia</i>	Nana
Salvia	<i>Salvia farinacea</i>	Rhea
Sedum	<i>Sedum spurium</i>	Dragon Blood Red Red Carpet Yellow
Spiraea	<i>Astilbe X arendsii</i>	Fanall
Spiraea	<i>Spiraea spp.</i>	Anthony Waterer Red Dolchica Froebeli Pink Goldenflame Red Snowmound White Van Houttee White
Spiraea, Garland	<i>Spiraea spp.</i>	
Spruce, Black Hills	<i>Picea glauca</i>	
Spruce, Colorado Blue	<i>Picea pungens</i>	
Spruce, Norway	<i>Picea abies</i>	
Spruce, White	<i>Picea glauca</i>	
Sourwood	<i>Oxydendrum arboreum</i>	
Sweetflag, Grassyy-Le		
Sweetgum	<i>Liquidambar styraciflua</i>	
Sycamore	<i>Platanus occidentalis</i>	
Tree Fern (Tiki Fern)	<i>Asparagus virgatus</i>	
Trumpet Flower, Evening	<i>Gelsemium sempervirens</i>	
Tulip	<i>Tulip, spp</i>	Apeldoorn
Verbena, Shrub	<i>Lantana sellowiana</i>	
Vervain	<i>Verbena spp.</i>	St. Paul
Viburnum	<i>Viburnum spp.</i>	American Cranberry Bush Arrowood Common Snowball European Cranberry Bush Linden

		Mohican Wright
Windmill Palm	<i>Trachycarpus fortunei</i>	
Xylosma	<i>Xylosma congestum</i>	
Yaupon	<i>Ilex vomitoria</i>	Dwarf
Yew	<i>Taxus cuspidata</i> <i>Taxus X media</i>	Denisiformis

*Ornamental species only: DO NOT USE ON FOOD PRODUCING TREES AND ORNAMENTALS

WEEDS CONTROLLED

When used as directed, this product will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence. This product will not control established grasses except for crabgrass in early stages of development, and broadleaf weeds, therefore, the area to be treated should be free of weeds prior to application.

GRASSES

barley (*Hordeum* spp.)
barnyardgrass (*Echinochloa crus-galli*)
bluegrass, annual (*Poa annua*)
brome (*Bromus* spp.)
crabgrass, large (*Digitaria sanguinalis*)
crabgrass, smooth (*Digitaria ischaemum*)
crabgrass, Southern (*Digitaria ciliaris*)
crowfootgrass (*Dactyloctenium aegyptium*)
dallisgrass (seedling) (*Paspalum dilatatum*)
goosegrass (*Eleusine indica*)
foxtail, green (*Setaria verdi*)
foxtail, yellow (*Setaria faberi*)
kikuyugrass (*Pennisetum clandestinum*)
oats, wild (*Avena fatua*)
ryegrass (annual & pernnial) (*Lolium* spp.)
smutgrass (*Sporobolus poireti*)

BROADLEAF WEEDS

bittercress (*Cardamine* spp.)
carpetweed (*Mollugo verticillata*)
chickweed (*Stellaria* spp.)
geranium, Carolina (*Geranium carolinianum*)
henbit (*Lamium* spp.)
knotweed, prostrate (*Polygonum saviculare*)
lespedeza, common (*Lespedeza striata*)
marestail (*Hippuris vulgaris*)
medic, black (*Medicago lupulina*)
mustard (*Brassica* spp.)
oxalis, buttercup (*Oxalis pes-caprae*)
pineappleweed (*Matricaria matricarioides*)
pigweed, redroot (*Amaranthus retroflexus*)
parsley-piert (*Alchemilla avrensis*)
purslane, common (*Portulaca oleracea*)
rocker, London (*Sisymbrium irio*)
shepherdspurse (*Capsella bursa-pastoris*)
speedwell, corn (*Veronica arvensis*)
spurge, garden (*Euphorbia hirta*)
spurge, prostrate (*Euphorbia humistrata*)
spurge, spotted (*Euphorbia maculata*)
woodsorrel, creeping (*Oxalis corniculata*)
woodsorrel, yellow (*Oxalis stricta*)

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Manufactured by:
The Andersons, Lawn Fertilizer Division, Inc.
P.O. Box 119
Maumee, OH 43537

Date Printed	6/2/15
Date Approved	5/8/03
Supersedes	5/26/98
Label Changes	Amendment (language for states with N and P restrictions)
Filename	9198-121c
Label Code	Rvamd06022015

NOTIFICATION

9198-121

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

07/01/2015

9198-121H
Page 1 of 13

(FRONT PANEL)
HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

The Andersons Dimension Herbicide III

Optional marketing claims:

- A Herbicide Plus Fertilizer Combination Product that Provides Control of Crabgrass and Other Weeds in Established Lawns and Ornamental Turfgrasses.
- Premium Granular Fertilizer that Feeds Your Lawn - Providing The Nutrients Needed for A Beautiful, Green and Vigorous Lawn.
- Contains Dimension® Herbicide - A Herbicide for Crabgrass Control!
- Can Be Applied Up to 4 Weeks Later Than Other Crabgrass Preventers and Controls Crabgrass All Season Long.
- For use on Lawns
- Controls Crabgrass: Pre-emergence & early Post-emergence
- Effective for up to 4 weeks after Crabgrass sprouts
- Use on cool-season & warm-season grasses
- Controls and/or suppresses over 10 broadleaf weeds
- Control Crabgrass in the Spring

ACTIVE INGREDIENTS:

Dithiopyr, 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S- dimethyl ester.....	0.164%
---	--------

INERT INGREDIENTS.....	<u>99.836%</u>
TOTAL.....	100.000%

® Dimension is a registered trademark of Dow AgroSciences LLC

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT _____ LBS.
Covers {over} X,XXX square feet

EPA Reg. No. 9198-121
EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
Underlined letter is first letter used in run code on bag

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

The Andersons Dimension Herbicide III

GUARANTEED ANALYSIS

Total Nitrogen (N).....	16%
Available Phosphate (P ₂ O ₅).....	4%
Soluble Potash (K ₂ O).....	4%
Primary Plant Food derived from: Urea, Muriate of Potash, Ammoniated Phosphates.	

For products being sold into Florida:

Florida Applications:

We recommend that you follow the Florida Green Industries BMP's at:

http://www.flaes.org/pdf/BMP_Book_final.pdf

We recommend that you follow the Florida Golf Course BMP's at: <http://www.flaes.org/pdf/gfbbmp07.pdf>

Nitrogen Notice: For those states with nitrogen application restrictions, including but not limited to MD, NH, NJ, and VA, do not apply more than x.xx lbs. of this product per 1,000 sq. ft.

Phosphate Notice: This lawn/turf fertilizer contains phosphorous and is only for nonagricultural use on (i) turf during its first growing season, (ii) on turf areas being repaired or renovated, and (iii) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not for routine maintenance of turf.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN

CAUTION

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply when weather conditions favor drift from treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 7.0 pounds per 1000 square feet (equal to 0.5 pound active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

Use only according to label instructions.

GENERAL INFORMATION

This product is a combination product that provides both crabgrass control and the required nutrients for optimum turfgrass growth all in a single product.

This product contains Dimension, a crabgrass preventer that can be applied up to 4 weeks later than other crabgrass preventers and provide up to 4 months control of crabgrass and other grass and broadleaf weeds prior to their establishment in lawns and ornamental turfgrasses. This product should be applied to lawns and ornamental turfgrasses that are well established (See "Use Precautions" section below for a list of the tolerant turfgrasses). This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

This product should only be applied to lawns and ornamental turfgrasses that are well established. Use of this product on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury.

The Andersons Dimension Herbicide III
can be applied to the following turfgrasses when applied as directed:

Cool-Season Turfgrasses

- Creeping Bentgrass
- Fine Fescues
- Kentucky Bluegrass
- Perennial Ryegrass
- Tall Fescues

Warm-Season Turfgrasses

- Bahiagrass
- Bermudagrass
- Buffalograss
- Carpetgrass
- Centipedegrass
- St. Augustinegrass
- Zoysiagrass

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

This product should be applied in the spring or fall prior to the weed seed germination in established lawns and ornamental turfgrasses.

This product can be applied up to 4 weeks later than other crabgrass preventers and still provide season-long weed control of crabgrass. Dimension herbicide provides preemergence control of crabgrass and other weeds when applied prior to their germination, and it also provides postemergence control of crabgrass. The postemergence control provided by this product is limited to crabgrass only and will provide control of crabgrass when applied up to 4 weeks after crabgrass germination. Application of this product later than 4 weeks after crabgrass germination will result in reduced control of crabgrass.

For control of other grass and broadleaf weeds, this product must be applied as a preemergent treatment. Only crabgrass can be controlled with delayed applications after germination.

Tips for Improved Weed Control: For best results, apply this product within a few days after mowing and delay further mowings after the application of this product until the lawn or ornamental turfgrass has been watered or received significant rainfall to dissolve the fertilizer granules. Improved weed control may be observed when treated lawn or ornamental turfgrass is watered or receives rainfall within a few days after application of this product.

Reseeding, Overseeding, or Sprigging of turfgrasses into areas that have been previously treated with this product should be delayed until 16 weeks after application. Reseeding, overseeding, or sprigging upto 16 weeks after application may prevent the establishment of the newly reseeded, overseeded or sprigged turfgrasses.

- DO NOT apply more than 7 pounds of this product per 1,000 square feet of turfgrass area during the year (1,000 square feet area = 100 feet x 10 feet).
- DO NOT apply to flowers, vegetables, or fruit trees.
- DO NOT use clippings from treated turf for mulching around flowers, vegetables or fruit trees.
- DO NOT apply this product to pastures
- Keep people and pets off treated areas until dust has settled.

APPLICATION DIRECTIONS

The Andersons Dimension Herbicide III will provide control of crabgrass and other grasses and broadleaf weeds listed below when applied prior to weed seed germination in established lawns and ornamental turfgrasses.

Weeds Controlled and/or Suppressed by The Andersons Dimension Herbicide III

Grass Weeds:

Annual Bluegrass
Annual Ryegrass
Barnyardgrass
Crabgrass
Crowfootgrass
Goosegrass

Green Foxtail
Kikuyugrass (seedling)
Perennial Ryegrass
Smutgrass (seedling)
Yellow Foxtail

Broadleaf Weeds:

Bittercress
Buttercup Oxalis
Carolina Geranium
Chickweed
Common Lespedeza
Common Purslane
Corn Speedwell

Creeping Oxalis
Henbit
Parsley-piert
Pineappleweed
Spotted Spurge
Yellow Woodsorrel

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

Application Timings:

This product may be applied either in the spring or in the fall for the control of weeds.

When this product is applied in the fall, a sequential or second application should be made the following spring to provide season-long control of weeds.

Length of Control:

This product will provide up to 4 months of weed control if applied according to label directions.

The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States that this product is being applied.

Please check the map below to determine which "Region" of the U.S. that applies to your particular use area.

Use "Regions" within the United States:



Recommended Use Rates:

Recommended Use Rates for The Andersons Dimension Herbicide III		
(Pounds of this product to apply per 1,000 square feet of turfgrass)		
<u>Region of U.S.</u>	<u>Length of Weed Control</u>	
	<u>2-3 Months Control</u>	<u>3-4 Months Control</u>
North	1.8	2.5
South	2.7	3.5

SUGGESTED SPREADER SETTINGS

(Available at Final Printing)

Broadcast Spreaders: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of product will ensure complete coverage.

Drop Spreaders: First, apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

Important: It is important to follow the directions as indicated on this quality lawn care product as over-application may cause injury to the lawn. Measure the size of the area to be treated and do not over or under apply. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Calibrate spreader prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. Close spreader when filling, stopping or turning. Do not hand spread.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

STORAGE: Keep container closed and away from food, feedstuffs and domestic water supplies.

DISPOSAL:

If empty: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

GENERAL INFORMATION - LANDSCAPE ORNAMENTALS

This product provides preemergence control of listed annual grass and broadleaf weeds in landscape ornamentals. It may be applied to soil surfaces where ornamental plants are grown for aesthetic purposes in ornamental gardens, and residential properties.

Ornamental plants on this label have shown tolerance to this product when applied as directed. It is impossible, however, to evaluate tolerance for all ornamental plant species or varieties or under all possible growing conditions. Under your growing conditions, a limited area involving only a few plants should be treated for tolerance evaluation prior to large-scale applications.

USE RATES

Apply this product with a properly calibrated spreader which will assure accurate, even particle distribution. This product may be applied as a single application or as a split application. Refer to the following table for product recommendations when treating a small area.

DO NOT apply more than 7 pounds per 1,000 square feet per year. Do not apply in Nassau and Suffolk Counties of New York.

**Pounds of Product
per 1,000 square feet**
7

**Ounces of Product
per 100 Square Feet**
11.2

APPLICATION TIMING AND RECOMMENDATIONS

This product is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination or to bare ground. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. Once an application of this product has been made, do not disturb the soil surface as the herbicide barrier will be broken.

PRECAUTIONS

- Only apply this product to established ornamentals.
- Do not apply this product directly to bare root of ornamental plants as injury may result.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply to soil around ornamental plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes or disease.

(BACK PANEL)
HOMEOWNER USE ONLY*(The above is for internal use only and will not appear on the final label copy)*

- To avoid foliar damage to ornamentals, irrigate immediately after application to remove herbicide particles from the foliage.
- Do not apply under conditions that would affect product distribution. Wind could affect distribution and move product onto sensitive species. Uneven product distribution will result in uneven weed control.
- Do not apply this product on grasses grown for seed.
- Do not graze livestock or feed forage cut from areas treated with this product.

TOLERANT ORNAMENTALS

When applied as directed under the conditions described on this label, ornamentals listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing condition. Follow directions given above to determine plant tolerance under your growing conditions prior to large scale use.

COMMON NAME	BOTANICAL NAME	TOLERANT CULTIVARS
Abelia, Dwarf	<i>Abelia X Grandiflora</i>	Nana
Ajuga	<i>Ajuga reptans</i> <i>Ajuga genevensis</i>	Bronze Bronze Beauty
Almond, Flowering	<i>Prunus gladiosa</i>	
Apple*	<i>Malus pumila</i>	
Arbovitae	<i>Thuja occidentalis</i>	Nigra Pyramidalis Smaragh Techny Woodwardii
Arborvitae, Golden	<i>Thuja orientalis</i>	
Aster, Chinese	<i>Callistephus chinensis</i>	Dwarf Queen
Ash, Green	<i>Fraxinus pennsylvanica</i>	
Ash, Mountain	<i>Sorbus acucparia</i>	
Ash, Purple	<i>Fraxinus americana</i>	
Azalea	<i>Rhododendron spp.</i>	Brillant Buccaneer Carror Chimes (Belgian) Exbury Fashion Hardijzer Beauty Hershey Red Higasa Hinocrimson Holland (Hybrid) Marion Lee Northern Lights Orange Cup Orchid Lights Snow Southern Charm
Azalea, Flame	<i>Rhododendron calendulaceum</i>	
Azalea, Kirishima		
Bamboo, Heavenly		
Barberry	<i>Berberis thunbergii</i>	Aurea Dwarf Pigmy Green Kobold Pygmy Red

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

		Rose Glow
Barberry, Purple		Atropurpurea
Bearberry (common)	<i>Arctostaphylos uva-ursti</i>	Massachusetts
Birch, River	<i>Betula nigra</i>	
Blueberry*	<i>Vaccinium spp.</i>	Bluecorp Blue Jay Jersey North Blue Northland
Bottlebrush	<i>Callistemon citrinum</i>	
Boxwood, Japanese		Japonica
Boxwood, Weller	<i>Buxus sempervirens</i>	
Broom	<i>Cytisus spp.</i>	Moonlight
Broom	<i>Genista pilosa</i>	Vancouver Gold
Bugle Carpet		
Carex, Variegated	<i>Carex</i>	
Camellia	<i>Camellia japonica</i> <i>Camellia sasangua</i>	Debutante Mathotiana Supreme Chansonette
Cedar, Red	<i>Juniperus virginiana</i>	
Cockscomb, Plumosa	<i>Celosia cristata</i>	Red Kewpie
Coleus	<i>Coleus blumei</i>	
Copper leaf	<i>Acalypha wilkesiana</i>	
Cotoneaster	<i>Cotoneaster apiculatus</i>	
Cycads	<i>Cycads revoluta</i>	
Coyotebrush	<i>Baccharis pelularis</i>	
Cypress, Bald	<i>Taxodium distichum</i>	
Cypress, Italian	<i>Cupressus sempervirens</i>	Glauca
Cypress, Leyland	<i>Cupressocyparis leylandii</i>	
Cypress, Japanese False	<i>Chamaecyparis obtusa</i>	Gracilis
Daffodil	<i>Narcissus spp.</i>	King Alfred
Daylily	<i>Hemerocallis spp.</i>	Aztec Gold Bright Yellow (Hybrid) Single Gold (Evergreen) Wilson's Yellow
Dogwood	<i>Cornus florida</i>	
Dogwood, American	<i>Cornus sericea</i>	
Douglas Fir	<i>Pseudotsuga menziesii</i>	
Dusty Miller	<i>Senecio cineraria</i>	
Elm, Drake	<i>Ulmus parvifolia</i>	
Euonymus	<i>Euonymus fortunei</i>	Argenteo-variegata Auereo-marginata Colorata Emerald Gaiety Emerald 'N Gold Gold Edge Gold Princess Silver King Tricolor Vegetus
Fan Palm, European	<i>Chamaerops humilis</i>	
Fan Palm, Mexican	<i>Washingtonia robusta</i>	
Fescue	<i>Festuca glauca</i>	

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

Fetterbush	<i>Leucothoe fontiana</i>	Rainbow
Ficus	<i>Ficus retusa</i>	Nitidia
Forsythia	<i>Forsythia X intermedia</i> <i>Spring Glory</i>	Arnold Dwarf Bronxensis Dwarf Lynwood Gold Meadowlark Weeping
Fir Fraser	<i>Abies fraseri</i>	
Gardenia	<i>Gardenia jasminoides</i>	Mystery Radicans
Geranium	<i>Pelargonium X hortorum</i>	
Gum	<i>Eucalyptus citriodora</i>	
Hawthorn	<i>Crataegus spp.</i>	Cockspur White Crimson Cloud Enchantress Jack Evans Washington White
Heather, Twisted	<i>Erica cinerea</i>	Mediterranean Pink
Hibiscus	<i>Hibiscus spp.</i>	Brilliant Hula Girl Blue Bird
Hemlock, Canada	<i>Tsuga canadensis</i>	
Holly	<i>Ilex spp.</i> <i>Ilex X meserveae</i> <i>Ilex X attenuata</i>	Blue Boy Blue Girl Burfodii China Girl Ciompacta Forsteri Hellerie Japanese Northern Beauty Needlepoint Nellie R. Stevens Savanah
Holly, Chinese	<i>Ilex cornuta</i>	
Holly, Japanese	<i>Ilex crenata</i>	
Holly, Yaupon	<i>Ilex vomitoria</i>	
Honeysuckle	<i>Lonicera japonica</i>	Clavey's Dwarf Halliana Tatarian Canadian White Zebelli Red Hosta
Hosta	<i>Hosta spp.</i> <i>Hosta lancifolia</i>	Algo Marginata
Ice Plant	<i>Carpobrotus edulis</i>	
Impatiens	<i>Impatiens wallerana</i>	
Iris	<i>Iris spp.</i>	Wedgewood Dwarf blue
Ivy, English	<i>Hedera helix</i>	Bulgaria Thorndale
Jasmine, Asian	<i>Trachelopermum asiaticum</i>	
Juniper	<i>Juniperus spp.</i>	Arcadia Armstrong Bar Harbor
	<i>Juniperus horizontalis</i>	Blue Chip Blue Pacific

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

		Blue Rug
	<i>Juniperus horizontalis</i>	Blue Star Broadmoor Buffalo Calgary Carpet Emerald Sea Emerald Spreader Endora Compacta Fuitlandi Green Gold Tip Hetzi Hughes Manhattan Blue Parsoni Pfitzeriana Plumosa Prince of Wales Procumbens Dwarf San Jose Sargent Blude Sargent Green Scandia Scopulorum Moonglow Scopulorum Skyrocket Spartan
	<i>Juniperus chinensis</i>	Tamariscifolia
	<i>Juniperus sabina</i>	Weberi Youngstown Yukon Belle
King Palm	<i>Archontophoenix cunninghamiana</i>	
Laurel, Australian	<i>Pittosporum tobira</i>	
Laurel, Mountain	<i>Kalmia latifolia</i>	
Leucothoe	<i>Leucothoe fontanesiana</i>	
Lily, Africian	<i>Agapanthus africanus</i>	Albus Peter Pan
Lily, African Blue		
Lily of the Valley	<i>Pieris japonica</i>	Mt. Fire
Lilyturf	<i>Liriope muscari</i>	Evergreen Giant Lilac Beauty Majestic Monroe White Variegata
Ligustrum, Japanese	<i>Ligustrum japonicum</i>	
Liriope, Green	<i>Liriope spicata</i>	
Maple, Japanese	<i>Acer japonicum</i>	
Maple, Norway	<i>Acer platanoides</i>	
Maple, Red	<i>Acer rubrum</i>	
Maple, Sivler	<i>Acer saccharinum</i>	
Maple, Sugar	<i>Acer saccharum</i>	
Marigold	<i>Tagetes patula</i>	Honeycomb Wheeler's Dwarf Variegata
Mock Orange*	<i>Philadelphus spp</i>	Golden

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

		Snowflake Double White
Mopndo Grass	<i>Ophiopogon japonicus</i>	
Moss Rose	<i>Portulaca grandiflora</i>	Sunnyside
Myrtle, Crape	<i>Lagerstoemia indica</i>	Faurei Langer Muskogee Standard Pink
Myrtle, Wax	<i>Myrica cerifer</i>	
Nadina	<i>Nadina domestica</i>	Compacta Nana
Oleander	<i>Nerium oleander</i>	Hardy Red Petite Pink Sister Agnes
Oak, Laurel	<i>Quercus laurifolia</i>	
Oak, Pin	<i>Quercus palustris</i>	
Oak, Red	<i>Quercus rubra</i>	
Oak, Southern L	<i>Quercus virginiana</i>	
Oak, Willow	<i>Quercus phellos</i>	
Oteospermum	<i>Osteospermum fruitcosum</i>	Wirligig
Palm, Bangalow		
Pampas Grass	<i>Cortadaria selloana</i>	
Paper Flower	<i>Bougainvillea glabra</i>	Barbara Karst
Pachysandra	<i>Pachysandra terminalis</i>	
Peach*	<i>Prunus persica</i>	
Periwinkle, Dwarf	<i>Vinca minor</i>	
Petunia	<i>Petunia X hybrida</i>	Picoti
Photinia, Red Tip	<i>Photinia X fraseri</i>	
Pieris	<i>Pieris japonica</i>	
Pine, Australian	<i>Pinus nigra</i>	
Pine, Japanese Black	<i>Pinus thumbergiana</i>	
Pine, Loblolly	<i>Pinus teada</i>	
Pine, Longleaf	<i>Pinus palustris</i>	
Pine, Mugho	<i>Pinus mugho</i>	
Pine, Scotch	<i>Pinus sylvestris</i>	
Pine, Slash	<i>Pinus eliottii</i>	
Pine, Swiss Mt.	<i>Pinus mugo</i>	
Pine, Virginia	<i>Pinus virginiana</i>	
Pine, White	<i>Pinus strobus</i>	
Pineapple, Guava	<i>Feijoa fellowaiana</i>	
Pittosporum, Japan		
Potentilla	<i>Potentilla fruticosa</i>	Abbotswood
Privet	<i>Ligustrum japonicum</i>	Golden Vicary Regal Texanum Yellow Tipped Wax
Privet, Glossy	<i>Ligustrum lucidum</i>	
Pyracantha	<i>Pyracantha koidzumii</i>	
Queen Pam	<i>Arecastrum rammanzoffianum</i>	
Quince, Japanese*		
Rhododendron	<i>Rhododendron spp</i>	Album Cunningham White Fashion Hardy

(BACK PANEL)
HOMEOWNER USE ONLY
(The above is for internal use only and will not appear on the final label copy)

		PJM Purple Gem Silvery Pink
Rhododendron, Carolina	<i>Rhododendron carolinianum</i>	
Rhododendron, Catawba	<i>Rhododendron catawbiense</i>	
Rodie Max (Rosebay)	<i>Rhododendron maximum</i>	
Rose	<i>Rosa banksiaae</i>	Luta
Rosemary*	<i>Rosmarinum officinalis</i>	
Rosemary, Bog	<i>Andromeda polifolia</i>	Nana
Salvia	<i>Salvia farinacea</i>	Rhea
Sedum	<i>Sedum spurium</i>	Dragon Blood Red Red Carpet Yellow
Spiraea	<i>Astilbe X arendsii</i>	Fanall
Spiraea	<i>Spiraea spp.</i>	Anthony Waterer Red Dolchica Froebeli Pink Goldenflame Red Snowmound White Van Houttee White
Spiraea, Garland	<i>Spiraea spp.</i>	
Spruce, Black Hills	<i>Picea glauca</i>	
Spruce, Colorado Blue	<i>Picea pungens</i>	
Spruce, Norway	<i>Picea abies</i>	
Spruce, White	<i>Picea glauca</i>	
Sourwood	<i>Oxydendrum arboreum</i>	
Sweetflag, Grassy-Le		
Sweetgum	<i>Liquidambar styraciflua</i>	
Sycamore	<i>Platanus occidentalis</i>	
Tree Fern (Tiki Fern)	<i>Asparagus virgatus</i>	
Trumpet Flower, Evening	<i>Gelsemium sempervirens</i>	
Tulip	<i>Tulip, spp</i>	Apeldoorn
Verbena, Shrub	<i>Lantana sellowiana</i>	
Vervain	<i>Verbena spp.</i>	St. Paul
Viburnum	<i>Viburnum spp.</i>	American Cranberry Bush Arrowwood Common Snowball European Cranberry Bush Linden Mohican Wright
Windmill Palm	<i>Trachycarpus fortunei</i>	
Xylosma	<i>Xylosma congestum</i>	
Yaupon	<i>Ilex vomitoria</i>	Dwarf
Yew	<i>Taxus cuspidata</i> <i>Taxus X media</i>	Deniseformis

***Ornamental species only: DO NOT USE ON FOOD PRODUCING TREES AND ORNAMENTALS**

WEEDS CONTROLLED

When used as directed, this product will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence. This product will not control established grasses except for crabgrass in early stages of development, and broadleaf weeds, therefore, the area to be treated should be free of weeds prior to application.

(BACK PANEL)
HOMEOWNER USE ONLY

(The above is for internal use only and will not appear on the final label copy)

GRASSES

barley (*Hordeum* spp.)
barnyardgrass (*Echinochloa crus-galli*)
bluegrass, annual (*Poa annua*)
brome (*Bromus* spp.)
crabgrass, large (*Digitaria sanguinalis*)
crabgrass, smooth (*Digitaria ischaemum*)
crabgrass, Southern (*Digitaria ciliaris*)
crowfootgrass (*Dactyloctenium aegyptium*)
dallisgrass (seedling) (*Paspalum dilatatum*)
goosegrass (*Eleusine indica*)
foxtail, green (*Setaria verdi*)
foxtail, yellow (*Setaria faberi*)
kikuyugrass (*Pennisetum clandestinum*)
oats, wild (*Avena fatua*)
ryegrass (annual & perennial) (*Lolium* spp.)
smutgrass (*Sporobolus poireti*)

BROADLEAF WEEDS

bittercress (*Cardamine* spp.)
carpetweed (*Mollugo verticillata*)
chickweed (*Stellaria* spp.)
geranium, Carolina (*Geranium carolinianum*)
henbit (*Lamium* spp.)
knotweed, prostrate (*Polygonum saviculare*)
lespedeza, common (*Lespedeza striata*)
marestail (*Hippuris vulgaris*)
medic, black (*Medicago lupulina*)
mustard (*Brassica* spp.)
oxalis, buttercup (*Oxalis pes-caprae*)
pineappleweed (*Matricaria matricarioides*)
pigweed, redroot (*Amaranthus retroflexus*)
parsley-piert (*Alchemilla avrensis*)
purslane, common (*Portulaca oleracea*)
rocker, London (*Sisymbrium irio*)
shepherdspurse (*Capsella bursa-pastoris*)
speedwell, corn (*Veronica arvensis*)
spurge, garden (*Euphorbia hirta*)
spurge, prostrate (*Euphorbia humistrata*)
spurge, spotted (*Euphorbia maculata*)
woodsorrel, creeping (*Oxalis corniculata*)
woodsorrel, yellow (*Oxalis stricta*)

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Manufactured by:
The Andersons, Lawn Fertilizer Division, Inc.
P.O. Box 119
Maumee, OH 43537

Date Printed	6/2/15
Date Approved	5/8/03
Supersedes	5/26/98
Label Changes	Amendment (language for states with N & P restrictions)
Filename	9198-121H
Label Code	Rvamd06022015